

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>C11D 9/04, 9/00, 17/00, 9/22</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 98/00505</b> <b>(43) International Publication Date:</b> 8 January 1998 (08.01.98)
<b>(21) International Application Number:</b> PCT/US97/11036 <b>(22) International Filing Date:</b> 25 June 1997 (25.06.97)  <b>(30) Priority Data:</b> 08/673,869      2 July 1996 (02.07.96)      US  <b>(71) Applicant (for all designated States except US):</b> JOHNSON & JOHNSON CONSUMER PRODUCTS, INC. [US/US]; Grandview Road, Skillman, NJ 08558 (US).  <b>(72) Inventor; and</b> <b>(75) Inventor/Applicant (for US only):</b> LAMBINO, Danilo, L. [PH/PH]; 135 K6h, East Kamias, Quezon City 1102 (PH).  <b>(74) Agents:</b> CIAMPORCERO, Audley, A., Jr. et al.; Johnson & Johnson, One Johnson & Johnson Plaza, New Brunswick, NJ 08933-7003 (US).		<b>(81) Designated States:</b> AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
<b>(54) Title:</b> CLEAR, COLORLESS SOAP BAR WITH SUPERIOR MILDNESS, LATHERING AND DISCOLORATION RESISTANCE		
<b>(57) Abstract</b>  The present invention relates to a clear colorless soap bar with superior mildness, lathering and discoloration resistance. The clear colorless soap consists of a blend of C12-C18 fatty acids neutralized with sodium hydroxide (NaOH) and triethanolamine (TEA). Excess TEA acts as a co-solvent and is responsible for clarity of the soap bar. The bar also contains a branched chain acid such as isostearic acid to break up crystallinity and add to product clarity. Low color and color stability are obtained by removing unsaturated fatty acids, and by the use of antioxidants (BHT and Vitamin E). Low levels of C6 to C10 fatty acids provide exceptional mildness.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

-1-

**CLEAR, COLORLESS SOAP BAR WITH SUPERIOR MILDNESS,  
LATHERING AND DISCOLORATION RESISTANCE**

**Field of the Invention**

5        This invention relates to a clear soap bar with exceptional resistance to discoloration on aging and which is extremely mild to the skin.

**Background of the Invention**

10        U.S. Patent 2,820,768 (Fromont) discloses that transparent soap can be made by mixing and heating to a temperature of 100 to 120°C a transparent alkali metal soap with the reaction product of a fatty acid containing no less than 18 carbon atoms with excess  
15 triethanolamine. The soap preferably contains approximately 30% castor oil to improve transparency. Ricinoleates obtained from castor oil were found to dissolve fatty acid salts such as stearates, thus inhibiting crystallization of the soap on cooling.

20        U.S. Patent 3,793,214 (O'Neill et. al.) discloses that transparent soap bars may be made by neutralizing a mixture of saturated fatty acids and C5 to C18 branched chain fatty acids with a neutralizing agent comprising a  
25 sodium compound and an alkanolamine, preferably, triethanolamine. The neutralizing compound contains sufficient sodium compound to neutralize at least 40 percent of the fatty acids. The neutralizing agent also contains sufficient alkanolamine to provide 15 to 45  
30 weight percent of free alkanolamine in the final soap bar composition. The soap contains from 10 to 20 parts of branched chain fatty acid for each 100 parts of soap. Examples of suitable branched chain acids are trialkyl

-2-

- acetic acids commonly known as neo-acids, and 2-ethylhexanoic acid. In the preparation of the bars disclosed, the fatty acids are heated with sodium hydrosulfite to a temperature of 130 to 210°F (54 to 99°C) with stirring until homogeneous, and to this are added a pre-blend of the neutralizing agent and water. O'Neill et. al. also disclose that other components, e.g., preservatives, antioxidants, colorants and perfumes may also be present in the formulation.
- Following the neutralization step, the other components are then added, the mixture is stirred until homogeneous, and it is then poured into molds to cool and form soap bars.
- U.S. Patent 4,290,904 (Poper et. al.) discloses that a transparent soap may be made by saponifying a fatty oil, preferably, a mixture of tallow, coconut and castor oils, with caustic soda, water and a polyhydric alcohol. The soap also contains a tetrakis (hydroxyalkyl) ethylenediamine, which may be added either before or after saponification. Additional surfactants to increase foaming and to stabilize the foam, such as amine oxides and alkyl diethanolamides, are desirably added. Other components that may be added include chelating agents, colors, antioxidants and perfumes.
- U.S. Patent 4,468,338 (Lindberg) discloses that a transparent soap bar that does not lose its transparency or otherwise darken over time can be formulated from a mixture of alkali metal and triethanolamine salts of C6 to C18 fatty acids, citric acid or one of its alkali

-3-

metal salts, an alkali metal metabisulfite and water.  
The fatty acids have an iodine number between 8 and 15.

U.S. Patent 4,758,370 (Jungermann, et. al.)  
5 discloses a process for the continuous production of  
transparent soap. In their process, a mixture of fatty  
acids, which may include coco fatty acids, stearic acid,  
oleic acid, ricinoleic acid and other acids, is blended  
in a first storage tank. Sodium hydroxide solution,  
10 which may or may not contain other agents such as  
triethanolamine, is maintained in a second storage tank.  
The contents of the two tanks are blended together in a  
heated, stirred tank reactor at precise stoichiometric  
ratios to produce a composition which when placed in  
15 molds and cooled, hardens to a transparent soap bar.  
Jungermann et al. claim the benefit of speed, easier  
control, and the production of a lighter, more uniform  
product. They also disclose that inclusion of materials  
such as antioxidants, BHA, BHT, tocopherol, tocopherol  
20 acetate, sodium metabisulfite, chelating agents EDTA and  
DTPA, isostearic acid, and neo-decanoic acid may be  
added without adversely affecting the primary  
characteristics required.

25 U.S. Patent 5,310,495 and European Patent 0335026B1  
(both to Hill, et. al.) disclose transparent soap bars  
made from carefully controlled compositions. These  
patents disclose compositions which comprise a mixture  
of alkanolammonium and alkali metal C12 to C22 fatty  
30 acid salts, the mole ratio of these being from 0.1 to  
less than 1.0. A liquid solvent system comprising water  
and triethanolamine is also present, the weight ratio of

-4-

these being from greater than 0.25 to less than 1.0.  
The weight ratio of total fatty acid salts to solvent  
must range from greater than 0.2 to less than 1.0.

While these patents suggest that unsaturated soaps give  
5 bars with a characteristic yellow color, they state that  
the soaps may contain unsaturation in accordance with  
commercially acceptable standards, though excessive  
unsaturation is normally avoided.

10 Philippine Utility Model 8018 (Dy Dumalasa, et.  
al.) teaches a transparent soap bar made from low iodine  
value fatty acids. Butylated hydroxy toluene (BHT) is  
added to the soap formulation to retard discoloration.  
The fatty acids used in this soap include 2-ethyl  
15 hexanoic acid, topped coconut fatty acids, triple  
pressed stearic acid and lauric acid.

Disadvantages of the soap bars produced according  
to the above disclosures is that the soaps exhibit  
20 varying degrees of clarity, color and color stability.

Furthermore, the soaps of the prior art tend to  
discolor to varying degrees on aging. This  
discoloration is especially pronounced at the higher  
temperatures encountered in warm climates where air  
25 conditioning is less common. Furthermore, the soap bars  
of the prior art tend to irritate sensitive skin.

Accordingly, it is an object of the present  
invention to provide a soap bar of exceptional clarity  
30 and low color.

-5-

A further object of the present invention to provide a clear, colorless soap bar which resists discoloration on aging.

- 5 Yet a further object of this invention to provide a soap bar that can be made from fatty acids that are entirely plant-derived.

- 10 In another object of the present invention there is provided a clear, colorless soap bar which is sufficiently mild to the skin to permit its regular use by individuals with sensitive skin, for example, infants.

15 **Summary of the Invention**

- A clear soap bar which resists discoloration on aging and which is extremely mild to the skin comprises 0.1 to 1.0 percent by weight of a reducing agent, about 40 to about 65 percent by weight of a mixture of alkali metal and alkanolamine salts of C6 to C22 carboxylic acids, about 35 to about 60 percent by weight of a solvent mixture comprising water and alkanolamine, and a discoloration-retarding effective amount of an antioxidant to retard the discoloration of the soap bar on aging, wherein about 2 to about 20 weight percent of the carboxylic acids are branched carboxylic acids, and wherein the carboxylic acids from which the carboxylic acid salts in the soap bar are derived have an iodine number less than or equal to about 2.0.

30

In a preferred embodiment, the reducing agent is selected from alkali metal metabisulfite, alkali metal

-6-

sulfite, alkali metal bisulfite and alkali metal hydrosulfite, and is most preferably sodium hydrosulfite. The reducing agent is present in the soap bar at a concentration of about 0.2 to about 0.6 percent by weight and most preferably at a concentration of 0.4 percent by weight. The alkanolamine is triethanolamine, about 50 to about 55 percent by weight of the soap bar comprises a mixture of sodium and triethanolamine salts of C6 to C22 carboxylic acids, the weight ratio of sodium salts to triethanolamine salts being preferably from about 25:75 to about 75:25, most preferably from about 45:55 to about 55:45; at least about 80 percent by weight and most preferably at least about 90 percent by weight of the carboxylic acids from which the carboxylic acid salts in the soap bar are derived are C12 to C18 straight chain, saturated carboxylic acids, the soap bar comprises from about 35 to 60 percent by weight and most preferably from about 40 to about 55 percent by weight of a solvent mixture which comprises from about 65 to about 85 percent and most preferably from about 70 to about 75 percent by weight triethanolamine. The antioxidant is selected from alkylated phenols and their derivatives, Vitamin E and its derivatives, and mixtures thereof, and more preferably, the antioxidant comprises a first antioxidant selected from butylated hydroxy toluene and butylated hydroxy anisole at a concentration of about 0.001 to about 0.1 percent by weight and a second antioxidant selected from Vitamin E and Vitamin E acetate at a concentration of about 0.01 to about 1.0 percent by weight. Even more preferably, the first antioxidant is butylated hydroxy toluene and is present in the soap bar at a concentration of about 0.005 to



-7-

about 0.05 percent by weight, and the second antioxidant is Vitamin E and is present in the soap bar at a concentration from about 0.05 to about 0.5 percent by weight. Further, more preferably the branched  
5 carboxylic acids are selected from the iso acids, neo acids, 2-ethyl hexanoic acid and mixtures thereof, and more preferably, the branched carboxylic acid is selected from isostearic acid, 2-ethyl-hexanoic acid and mixtures thereof and comprises from about 2 to about 10  
10 percent of the carboxylic acids from which the soap bar is derived.

The soap bar may also contain ancillary agents such as foam stabilizers, humectants, emollients, fragrances  
15 and chelating agents. Examples of foam stabilizers include alkyl monoethanolamides, alkyl diethanolamides, acyl sarcosinates, acyl taurates, acyl isethionates, acyl lactates, alkyl amine oxides, alkyl betaines and mixtures thereof. Examples of humectants include  
20 glycerine, propylene glycol, butylene glycol, polyethylene glycol, and mixtures thereof. Examples of emollients include mineral oil, vegetable oil, silicone oils, synthetic and semisynthetic emollient esters and mixtures thereof. Examples of chelating agents include  
25 the tetrasodium salt of ethylenediamine tetraacetic acid and the pentasodium salt of diethylenetriamine pentaacetic acid.

In the preferred soap bars of the present  
30 invention, the carboxylic acids from which the carboxylic acid salts in the soap bar are derived comprises from zero to less than about 0.5 percent by

-8-

weight of C6 to C10 linear, straight chain carboxylic acids. In some regions of the world, it is preferred that the carboxylic acids from which the soap bar is derived be entirely of plant origin, and this  
5 requirement may be accommodated by the soap bars of the present invention.

Also disclosed is a process for making the soap bar which includes the steps of providing a mixture of C6 to  
10 C22 carboxylic acids wherein about 2 to about 20 weight percent of the carboxylic acids are C8 to C18 branched carboxylic acids and wherein the carboxylic acids have an iodine number less than or equal to about 2.0, heating the mixture of C6 to C22 carboxylic acids to  
15 produce a molten mixture of carboxylic acids, admixing a reducing agent with the molten carboxylic acids, admixing less than a molar equivalent of alkali metal hydroxide with the C6 to C22 carboxylic acids, affording a mixture of unneutralized carboxylic acids with  
20 neutralized carboxylic acids, admixing alkanolamine with the mixture of unneutralized carboxylic acids and neutralized carboxylic acids, affording a mixture of alkali metal and alkanolamine salts of the C6 to C22 carboxylic acids, providing sufficient excess of  
25 alkanolamine to function as a solvent which increases the clarity of the soap bar, thereby providing a mixture of carboxylic acid salts and solvent comprising alkanolamine, admixing a discoloration-retarding effective amount of an antioxidant to the carboxylic  
30 acid salts and solvent, transferring the molten mixture of salts and solvent into molds, and cooling the molds to harden the molten soap into solid soap. In the

-9-

neutralization of the acids, the alkali metal hydroxide may be mixed with the acids before addition of the alkanolamine, the alkanolamine addition may precede the neutralization with alkali metal hydroxide, or the  
5 neutralization with alkali metal hydroxide and alkanolamine may be conducted simultaneously.

#### Brief Description of the Drawing

Figure 1 is a graph showing the effect of aging on the a\* color value of the soap bars of the present  
10 invention.

Figure 2 is a graph showing the effect of aging on the b\* color value of the soap bars of the present  
15 invention.

#### Detailed Description of the Invention

The present invention is a clear soap bar of exceptionally low color which resists discoloration on aging, and the process for making this soap bar. The  
20 soap bar of the present invention is very mild, and it can, therefore, be used on a regular basis by individuals with sensitive skin, for example, infants.

25 The soap bar of the present invention comprises the following:

- a. 0.1 to 1.0 percent by weight of a reducing agent,
- b. about 40 to about 65 percent by weight of a mixture of alkali metal and alkanolamine salts of C6 to C22  
30 carboxylic acids,
- c. about 35 to 60 percent by weight of a solvent mixture comprising water and an alkanolamine,

-10-

d. a discoloration-retarding effective amount of an antioxidant to retard the discoloration of the soap bar on aging.

5           In the soap bars of the present invention, about 2 to about 20 weight percent of the carboxylic acids from which the carboxylic acids salts are derived are branched chain carboxylic acids. The carboxylic acids possess very low levels of unsaturation, as evidenced by  
10           an iodine number less than or equal to a value of about 2.0.

          The soap bars of the present invention contain a reducing agent at a concentration of about 0.1 to about  
15           1.0 percent by weight. The reducing agent is preferably an inorganic sulfurous salt selected from alkali metal metabisulfite, alkali metal sulfite, alkali metal bisulfite and alkali metal hydrosulfite. The alkali metal in the inorganic sulfurous salt may be sodium or  
20           potassium, although sodium is preferred. The preferred reducing agent is sodium hydrosulfite, and it is most preferably added to the soap bar ingredients at a concentration of about 0.2 to about 0.6 percent by weight. The reducing agent is believed to function by  
25           reducing the color-bodies that are present in some of the fatty acid ingredients, as well as some of the impurities that contribute to the formation of color bodies in the soap over time.

30           The soap bars of the present invention contain about 40 to about 65 percent by weight of a mixture of alkali metal and alkanolamine salts of C6 to C22 fatty

-11-

acids. The alkali metal may be sodium, potassium, or mixtures of sodium and potassium, although sodium is preferred. The alkanolamine is preferably triethanolamine, although minor amounts of other alkanolamines such as diethanolamine may also be present.

The carboxylic acids salts in the soap bars of the present invention are formed by the neutralization of fatty acids with a mixture of alkaline materials comprising alkali metal hydroxide and an alkanolamine, preferably, sodium hydroxide and triethanolamine. Since sodium hydroxide is the stronger base, it will react preferentially with the fatty acids in the neutralization step. Thus, in order to form a mixture of alkali metal and alkanolamine salts, less than one mole of alkali metal hydroxide must be used for each mole of fatty acid. Any fatty acid left unneutralized by the molar deficiency of alkali metal hydroxide will then be neutralized by the alkanolamine.

The weight ratio of alkali metal carboxylic acid salt to alkanolamine salt in the soap bars of the present invention preferably ranges from 25:75 to about 75:25, and more preferably ranges from 55:45 to 45:55. For example, using the distribution of fatty acids described in the following Example 1, the carboxylic acids have an average molecular weight of about 236. In Example 1, the carboxylic acids are neutralized with sodium hydroxide and triethanolamine. The weight ratio of sodium salts of the carboxylic acids to the triethanolamine salts of the carboxylic acids in the

-12-

5 soap bar made according to Example 1 is 27.4:24.7, or 1.11:1. This ratio of salts is produced by reacting each mole of carboxylic acids with about 0.63 moles of sodium hydroxide and neutralizing the remainder of the acids with triethanolamine.

10 In forming the fatty acid salts comprising the soap bars of the present invention, the neutralization step may be conducted by sequentially reacting the fatty acids with each of the alkaline materials added separately, or by reacting the fatty acids with a combination of the alkaline materials.

15 The carboxylic acid salts which comprise the soap bars of the present invention are derived from C6 to C22 fatty acids. Preferably, at least about 80 percent by weight of the carboxylic acids should be linear, straight chain, saturated acids containing from 12 to 18 carbon atoms. The presence of unsaturation in the fatty acids has been found to contribute to the discoloration of the soap bars as initially formed as well as the further discoloration over time. Accordingly, the fatty acids used in the soap bars of the present invention should have a low degree of unsaturation. The degree of unsaturation in fatty acids is often indicated by the iodine number or iodine value, both of these terms used interchangeably herein. The iodine number may be determined by such methods as AOAC Official Method 920.158, which is incorporated herein by reference (Official Methods of Analysis of AOAC International, edited by Patricia Cunniff, Sixteenth Edition, 1995, Volume II, Chapter 41, page 6-7). The fatty acids from

20  
25  
30

-13-

which the present soap bars are derived preferably have an iodine number of less than about 5, and more preferably have an iodine number less than about 2.0

5           The presence of straight chain C6 to C10 acids or salts derived therefrom in the soap bars is believed to cause irritation of the skin. Accordingly, the carboxylic acids from which the soap bars of the present invention are derived preferably contain less than about  
10 0.5 percent by weight of C6 to C10 linear straight chain acids.

          Thus, the preferred fatty acids used in the soap bars of the present invention are low in unsaturated  
15 fatty acids and have a low concentration of C6 to C10 fatty acids. To accommodate both of these criteria, it is preferred to utilize purified single component or multiple component fatty acid fractions. Different acids can then be blended to optimize the final  
20 properties of the finished soap bar. The following are examples of fatty acids that are useful for preparing the soap bars of the present invention:

          Philacid 1200® (United Coconut Chemicals) contains  
25 at least 99 percent by weight lauric acid. It is obtained by saponifying crude coconut oil and distilling the crude fatty acids so obtained. The material has an iodine value of less than 0.3.

30           Philacid 1400® (United Coconut Chemicals) contains at least 99 percent by weight of myristic acid. It is also obtained by saponifying crude coconut oil and

-14-


distilling the crude fatty acids. This material also has an iodine value of less than 0.3.

Pristerene 4900® is a commercial low iodine value fatty acid from Unichema International, The Netherlands.

It is a mixture of fatty acids with the following typical composition:

4% lauric/myristic acids  
10 46% palmitic acid  
49% stearic acid  
1% oleic acid.

The carboxylic acids from which the soap bars of the present invention are derived comprise from about 2.0 to about 20 percent by weight of C8 to C18 branched chain carboxylic acids. The branched chain acids serve to break up the crystallinity of the carboxylic acid salts, thereby enhancing the clarity of the soap bar. Illustrative examples of branched chain acids that are useful in the soap bars of the present invention are trialkyl acetic acids, otherwise known as neo acids, of the formula:



25

wherein R, R' and R'' are all alkyl groups which may be the same or different. An example of a neo acid useful in the soap bars of the present invention is neodecanoic acid.

30



-15-

Other branched chain acids that are effective in the soap bars of the present invention are 2-ethyl hexanoic acid, and iso acids such as isostearic acid.

5 An example of an iso acid useful in the soap bars of the present invention is Prisorine 3505®, (Unichema International, The Netherlands), which has the following typical composition:

10 2.5% lauric/myristic acid, 10% branched C16 acids, 6% linear palmitic acid, 65% branched C18 acids, 2% linear stearic acid, 2.5% oleic acid, 8% branched C20 acids and 4% C22 acids. Thus, this material contains a total of about 83 weight percent branched carboxylic acids.

15

The soap bars of the present invention contain about 35 to about 60 percent, preferably about 40 to about 45 percent by weight of a solvent mixture. The solvent mixture comprises triethanolamine and water  
20 containing from about 65 to 85 percent, preferably about 70 to about 75 percent by weight of triethanolamine. Thus, in the preparation of the soap bar of the present invention, fatty acids are preferably neutralized with sodium hydroxide and triethanolamine, with sufficient  
25 excess of triethanolamine to provide the amount required for the solvent mixture. The presence of the solvent mixture in the soap bar further enhances the clarity of the obtained soap bar.

30 The soap bars of the present invention contain a discoloration-retarding effective amount of an

-16-

antioxidant to stabilize the discoloration of the bar with time. Suitable antioxidants include alkylated phenols such as butylated hydroxy toluene (BHT) and butylated hydroxy anisole (BHA), and Vitamin E (DL- $\alpha$ -tocopherol) and its derivatives such as Vitamin E acetate. A combination of antioxidants, such as BHT and Vitamin E, has been found to be especially effective at retarding the discoloration of the soap bars of the invention. For example, a soap bar preferably contains 0.001 to about 0.1 percent by weight of BHT and 0.01 to about 1.0 percent by weight of Vitamin E. More preferably, the bar contains from 0.005 to about 0.05 percent by weight of BHT and about 0.05 to about 0.5 percent by weight of Vitamin E.

15

The soap bars of the present invention may also contain suitable ancillary agents. Examples of such agents are foam stabilizers, humectants, emollients, chelating agents and fragrances. The foam stabilizers that may be useful in the soap bars of the present invention include alkyl monoethanolamides, alkyl diethanolamides, acyl sarcosinates, acyl taurates, acyl isethionates, acyl lactates, alkyl amine oxides, alkyl betaines and mixtures thereof. An example of a useful and effective foam stabilizer is cocamide DEA, the diethanolamide derived from coconut fatty acids with diethanolamine. This material may also be referred to as coco diethanolamide.

25

-17-

Humectants that may be useful in the soap bars of the present invention include glycerine, propylene glycol, butylene glycol, polyethylene glycol, and mixtures thereof. Glycerine is a preferred humectant in the soap bars of this invention. The presence of humectants in the soap bar leave the user with the feeling that the soap does not dry out the skin after use.

Emollients that may be useful in the soap bars of the invention include mineral oil, vegetable oil, silicone oils, synthetic and semisynthetic emollient esters and mixtures thereof. Mineral oil is a preferred emollient. The presence of emollients in the soap bar leaves the user's skin with a soft, silky feeling after use of the soap bar.

The presence of heavy metal ions is believed to catalyze reactions that contribute to discoloration of the soap. Accordingly, it is advantageous to include chelating agents in the soap bar formulation. Examples of chelating agents that may be used include the tetrasodium salt of ethylenediamine tetraacetic acid (EDTA) and the pentasodium salt of diethylenetriamine pentaacetic acid (Na<sub>5</sub>DTPA).

The following examples are illustrative of the soap bars of the present invention without intending to limit the invention in any manner. In the following examples, amounts of materials used are expressed in parts by

-18-

weight based on a total of 100 parts of material added to the soap bar formulation.

### Example 1

- 5        The following ingredients were charged to a stirred, jacketed Britannia reaction tank pre-heated to 65 to 70°C:

	<u>Ingredient</u>	<u>Chemical Name</u>	<u>Parts by weight</u>
10	Prisorene 3505®	Isostearic Acid	4
	Pristerene 4900®	Palmitic/Stearic Acid	17
	Philacid 1200®	Lauric Acid	13
	Philacid 1400®	Myristic Acid	6.3

- 15        The combined fatty acids have an iodine value of 0.6 and contain 0.2 percent by weight of C6 to C10 straight chain carboxylic acids.

- 20        Heating of the acid mixture was continued until its temperature is 60°C. 1.2 parts of a 33.3 percent by weight solution of sodium hydrosulfite was added, and the mixture was stirred for 10 minutes. 0.01 parts of BHT were then added. The acids were neutralized by the addition of 55.1 parts of a solution containing 7.7%  
25        sodium hydroxide, 75.5% triethanolamine and 16.8% water, the percentages of each of these components being percent by weight. The rate of addition was controlled to maintain the temperature of the acids between 78 and 80°C. The following ingredients were then added:

30

<u>Ingredient</u>	<u>Chemical Name</u>	<u>Parts by weight</u>
Versenex 80	Na5DTPA (    % in H <sub>2</sub> O)	0.5
Glycerine		1.0
mineral oil		0.5

-19-

coco diethanolamide	1.0
Fragrance --	0.4
Vitamin E	0.1

5           The molten soap was poured into molds and allowed to cool to room temperature. When the soap achieved the desired level of hardness, the soap was subjected to cutting, stamping and finishing operations.

10           The finished soap had the composition shown in the following Table 1:

-20-

Table 1

Component	Composition (percent by weight)
sodium soap	27.4
triethanolamine soap	24.7
water	12.0
triethanolamine	32
sodium hydrosulfite	0.4
BHT	0.01
Glycerine	1.0
mineral oil	0.5
coco diethanolamide	1.0
fragrance	0.4
pentasodium-diethylenetriaminepentaacetate	0.5
Vitamin E	0.1

The clarity of the soap was assessed by measuring the transmittance of 800 nm light through a 20 mm thick bar. The soap of example 1 had a percent transmittance of 52.3 when dry and 83.0 when wet.

#### Example 2

The method of Example 1 was repeated wherein the fragrance and Vitamin E were omitted.

#### Example 3

The method of Example 1 was repeated wherein Vitamin E was omitted.

-21-

Example 4

The method of Example 1 was repeated wherein isostearic acid was replaced by the same amount of 2-ethyl hexanoic acid, only 0.2 parts of fragrance was added and Vitamin E was omitted.

Examples 5 through 7

The method of Example 1 was repeated with the following amounts of Vitamin E added to the formulation:

10

<u>Example #</u>	<u>Amount of Vitamin E added</u> (weight %)
5	0.03
6	0.05
7	0.20

The ingredients in each of these examples is summarized in the following Table 2:

-22-

Table 2

Component	Ex. 1	Ex. 2	Ex. 3	Ex. 4	Ex. 5	Ex. 6	Ex. 7
Prisorene 3505®	4	4	4	0	4	4	4
2-ethyl hexanoic acid	0	0	0	4	0	0	0
Pristerene 4900®	17	17	17	17	17	17	17
Philacid 1200®	13	13	13	13	13	13	13
Philacid 1400®	6.3	6.4	6.3	6.3	6.3	6.3	6.3
sodium hydrosulfite	0.4	0.4	0.4	0.4	0.4	0.4	0.4
sodium hydroxide	4.2	4.3	4.2	4.3	4.2	4.2	4.2
water	10.0	10.1	10.1	10.1	10.0	10.0	10.0
triethanol- amine	41.6	41.8	41.6	41.8	41.6	41.6	41.6
Glycerine	1.0	1.0	1.0	1.0	1.0	1.0	1.0
mineral oil	0.5	0.5	0.5	0.5	0.5	0.5	0.5
coco diethanol- amide	1.0	1.0	1.0	1.0	1.0	1.0	1.0
pentasodium- diethylene- triaminepenta acetate	0.5	0.5	0.5	0.5	0.5	0.5	0.5
BHT	0.01	0.01	0.01	0.01	0.01	0.01	0.01
fragrance	0.4	0.0	0.0	0.0	N/A	N/A	N/A
Vitamin E	0.10	0.0	0.0	0.0	0.03	0.05	0.20

SUBSTITUTE SHEET (RULE 26)



-23-

The color of the soap bars immediately upon manufacture was assessed visually and compared against commercial clear bars. All of the bars made according to Examples 1 through 7 above appeared completely clear.

5 In contrast, commercially available bars ranged in color from faint yellow to dark brown.

One method to assess colors quantitatively is by use of CIE color space values. Aspects of this method are described in American Society of Testing and Materials (ASTM) Standard Method E-308-95 and in The United States Pharmacopeia (USP23)/The National Formulary (NF18) in Section 1061 entitled Color-Instrumental Measurement, both of these references  
10  
15 incorporated herein by reference. In this method, colors are assessed with the aid of a colorimeter. The software associated with the instrument produces three parameters, known as L\*, a\* and b\*, which are believed to correlate with the three separate stimuli with which  
20 colors are perceived by the human visual cortex. In this method, the L\* parameter measures the change in sample appearance along a gray scale, which ranges from 0 for theoretical black to 100 for white. The a\* and b\* parameters measure the red-versus-green and yellow-versus-blue attributes of the sample, respectively.  
25 Increases in the value of the a\* parameter correlate with increasing redness in the sample. Increases in the b\* parameter correlate with increasing yellow color in the sample.

30

The appearance of the as-produced bars was assessed colorimetrically by the above-indicated method using a

-24-

Hunter Associates Laboratory, Inc. Miniscan Portable Spectrocolorimeter. Colorimetric values were determined using bars of 20 mm thickness. The results are presented in Table 3 below:

5

-25-

Table 3<sup>1, 2</sup>

Bar	Appearance	Iodine Value	Transmittance <sup>1</sup> (% @ 800 nm)		L*	a*	b*
			Dry	Wet			
Example 1	clear, colorless	0.6	52.3	83.0	52.3	-1.2	2.2
Example 2	clear, colorless	0.6	N/A	N/A	52.2	-1.1	2.0
Commercial Bar A	faint yellow	36.7	33.8	75.7	52.0	0.3	8.1
Commercial Bar B	orange brown	82.8	26.4	60.3	37.8	5.0	3.6
Commercial Bar C	deep red	29.0	43.6	68.1	34.5	11.9	11.7

<sup>1</sup> Transmittance values measured using Perkin Elmer Lambda 2 UV/VIS Spectrophotometer on a bar thickness of 20 mm and at a wavelength of 800 nm.

5

<sup>2</sup> L\*, a\* and b\* values determined using a Hunter Associates Laboratory, Inc. Miniscan Portable Spectrocolorimeter.

-26 -

The transmittance data indicate the superior clarity of the soap bars of the present invention relative to commercially available clear soap bars.

5

Example 8

Samples produced by Examples 1, 3, 5, 6 and 7 were subjected to accelerated aging tests by heating to 49°C.

Samples were pulled at 1-week intervals, up to a period of 8 weeks. The results of these experiments are

10 summarized in the following Table 4:

Table 4			
Soap Bar of Example #	Amount on Vitamin E in soap bar (weight percent)	Number of weeks at 49°C	Visual Color
3	0	5	colorless
3	0	6	yellow
3	0	7	orange
3	0	8	dark red
5	0.03	7	colorless
5	0.03	8	orange
6	0.05	7	colorless
6	0.05	8	faint yellow
1	0.10	7	colorless
1	0.10	8	faint yellow
7	0.02	7	faint yellow
7	0.02	8	orange

From the visual appearance of the aged soap bars as reflected in the data in Table 4, it is evident that the color stability of the soap bar is enhanced at all levels of Vitamin E employed in these experiments. Furthermore, an optimum in stability exists at a Vitamin E concentration ranging from about 0.03 to about 0.2 percent by weight, and more preferably, from about 0.05 to about 0.10 percent by weight.

-27 -

The order of color stability of these bars, as evidenced by their color at week 8, is as follows, going from most stable to least stable:

5

Example 1 > Example 6 > Example 7 > Example 5 > Example 3

Figures 1 and 2 show the effect of various levels of Vitamin E on the a\* and b\* color values for the soap bars subjected to the above-described aging studies. Soap bars were pulled at one-week intervals and their color values were measured using the Hunter Associates Laboratory Miniscan Portable Spectrocolorimeter. Results for the soap bars that were subjected to these aging studies are shown in the following Table 5:

Table 5

Soap Bar of Example #	Amount on Vitamin E in soap bar (weight percent)	Symbol in Figures 1 and 2
3	0	diamond
5	0.03	square
6	0.05	triangle
1	0.10	X
7	0.02	*

20

As shown in Figure 1, even the lowest amount of Vitamin E used in the study, 0.03% had an effect on reducing the a\* color values of the soap bar in the accelerated aging test. Further levels of Vitamin E to 0.20% by weight had no further effect. The effect of Vitamin E on the b\* values is shown in Figure 2, which

25

-28 -

confirms an optimum level of about 0.05 to 0.10% by weight of Vitamin E in the formulation.

#### Example 9

5       The mildness of the soaps of the present invention was assessed using the Soap Chamber Test (P. J. Frosch and A. M. Kligman, *Am. Acad. Dermatol.*, 1:35-41, 1979). The test is designed to measure the irritancy of soaps using 5 consecutive weekday exposures to 8% solutions  
10 with readings of scaling, redness and fissuring on the following Monday. The test was conducted according to the following procedure: A Finn chamber on Scanpor tape containing a filter paper disc was used. 100  $\mu$ l of 8.0% soap solution was added by dropper onto the filter paper  
15 disc which was applied to the ventral skin of the forearm. On the first day, the fresh solutions were applied for 24 hours. On the next 4 days, the solutions were applied for 6 hours each day. The test site was evaluated on the Monday morning following the procedure  
20 using the following grading system:

#### Erythema

- 1+ Slight redness, spotty or diffuse
- 25 2+ Moderate, uniform redness
- 3+ Intense redness
- 4+ Fiery red with edema

#### Scaling

- 30 1+ Fine
- 2+ Moderate
- 3+ Severe with large flakes

#### Fissures

-29 -

- 1+ Fine cracks
- 2+ Single or multiple broader fissures
- 3+ Wide cracks with hemorrhage or exudation

5        The average of each of these parameters is calculated for all subjects (at least 20), and the values are summed to give a total score.

10        A soap bar made according to Example 4 was tested for irritancy using the above-described Soap Chamber Test. The total score for the soap bar of Example 4 was zero, indicating no detectable level of irritation. Although the presence of C6 to C10 straight chain acids is expected to contribute to irritation of the soap, the  
15        soap made according to Example 4, which contains 4% by weight of 2-ethyl hexanoic acid, a branched chain C8 acid, surprisingly shows no detectable levels of irritancy.

20

#### Example 10

Irritation values were measured for several commercial soaps using the method of Example 9, and the results were correlated against C6 to C10 fatty acid  
25        content as shown in Table 6 below:

Table 6

Soap Sample	C6 to C10 straight chain acid content (weight percent)	Irritation score
Example 4	0.2	0
Commercial Soap D	1.6	0.13
Commercial Soap E	2.3	0.65
Commercial Soap F	3.4	0.56

-30 -

As indicated in the table, the irritation score  
correlates with C6 to C10 straight chain acid content  
5 with a correlation coefficient of 0.89.



-31 -

I claim:

1. A clear soap bar comprising:
  - 5 (a) 0.1 to 1.0 percent by weight of a reducing agent;
  - (b) about 40 to about 65 percent by weight of a mixture of alkali metal salts of C6 to C22 carboxylic acids and alkanolamine salts of C6 to C22 carboxylic acids;
  - (c) about 35 to about 60 percent by weight of a solvent
  - 10 mixture comprising water and alkanolamine; and
  - (d) a discoloration-retarding effective amount of an antioxidant; wherein about 2 to about 20 weight percent of the carboxylic acids are C8 to C18 branched carboxylic acids, and wherein the carboxylic acids have
  - 15 an iodine number less than or equal to about 2.0.
2. The soap bar of claim 1 wherein the reducing agent is selected from the group consisting of alkali metal metabisulfite, alkali metal sulfite, alkali metal
- 20 bisulfite and alkali metal hydrosulfite.
3. The soap bar of claim 2 wherein the reducing agent is alkali metal hydrosulfite at a concentration of about 0.2 to about 0.6 percent by weight.
- 25 4. The soap bar of claim 3 wherein the alkali metal is sodium and the alkanolamine is triethanolamine.
5. The soap bar of claim 1 wherein at least about 80
- 30 percent by weight of the carboxylic acids from which the carboxylic acid salts in the soap bar are derived are C12 to C18 straight chain, saturated carboxylic acids.
6. The soap bar of claim 1 wherein the alkali metal
- 35 carboxylic acid salts and alkanolamine carboxylic acid

-32 -

salts are present in the soap bar in a weight ratio ranging from about 25:75 to about 75:25.

7. The soap bar of claim 1 wherein the alkanolamine is triethanolamine and the solvent mixture comprises from about 65 to about 85 percent by weight triethanolamine.

8. The soap bar of claim 1 wherein the antioxidant is selected from the group consisting of alkylated phenols and their derivatives, Vitamin E and its derivatives, and mixtures thereof.

9. The soap bar of claim 8 wherein the antioxidant comprises a first antioxidant selected from butylated hydroxy toluene and butylated hydroxy anisole at a concentration of about 0.001 to about 0.1 percent by weight and a second antioxidant selected from Vitamin E and Vitamin E acetate at a concentration of about 0.01 to about 1.0 percent by weight.

10. The soap bar of claim 9 wherein the first antioxidant is butylated hydroxy toluene and is present in the soap bar at a concentration of about 0.005 to about 0.05 percent by weight, and the second antioxidant is Vitamin E and is present in the soap bar at a concentration from about 0.05 to about 0.5 percent by weight.

11. The soap bar of claim 1 wherein the branched carboxylic acids are selected from the group consisting of iso acids, neo acids, 2-ethyl hexanoic acid and mixtures thereof.

-33 -

12. The soap bar of claim 11 wherein the branched carboxylic acids are selected from the group consisting of isostearic acid, 2-ethyl-hexanoic acid and mixtures thereof.

5

13. The soap bar of claim 1 which further comprises ancillary agents selected from the group consisting of foam stabilizers, humectants, emollients, fragrances and chelating agents.

10

14. The soap bar of claim 13 wherein the foam stabilizer is selected from the group consisting of alkyl monoethanolamides, alkyl diethanolamides, acyl sarcosinates, acyl taurates, acyl isethionates, acyl lactates, alkyl amine oxides, alkyl betaines and mixtures thereof.

15

15. The soap bar of claim 13 wherein the humectant is selected from the group consisting of glycerine, propylene glycol, butylene glycol, polyethylene glycol, and mixtures thereof.

20

16. The soap bar of claim 13 wherein the emollient is selected from the group consisting of mineral oil, vegetable oil, silicone oils, synthetic and semisynthetic emollient esters and mixtures thereof.

25

17. The soap bar of claim 13 wherein the chelating agent is selected from the tetrasodium salt of ethylenediamine tetraacetic acid and the pentasodium salt of diethylenetriamine pentaacetic acid.

30

18. The soap bar of claim 1 wherein the carboxylic acids from which the carboxylic acid salts in the soap bar are

-34 -

derived comprises from zero to less than about 0.5 percent by weight of C6 to C10 linear, straight chain carboxylic acids.

- 5 19. The soap bar of claim 1 wherein the C6 to C22 carboxylic acids are entirely of plant-derived origin.
20. The soap bar of claim 1 wherein:
- 10 (a) the reducing agent is sodium hydrosulfite, present at a concentration of about 0.2 to about 0.6 percent by weight;
- (b) the alkali metal is sodium and the alkanolamine is triethanolamine and the soap bar comprises from about 40 to about 65 percent by weight of sodium and
- 15 triethanolamine salts of C6 to C22 carboxylic acids;
- (c) at least about 80% by weight of the carboxylic acids from which the carboxylic acid salts in the soap bar are derived are C12 to C18 linear, straight chain, saturated carboxylic acids;
- 20 (d) the solvent mixture comprises water and triethanolamine, the solvent mixture comprising from about 65 to about 85 percent by weight triethanolamine;
- (e) the antioxidant comprises a first antioxidant and a second antioxidant, wherein the first antioxidant is
- 25 butylated hydroxy toluene and is present in the soap bar at a concentration of 0.005 to about 0.05 percent by weight, and the second antioxidant is Vitamin E and is present in the soap bar at a concentration from about 0.05 to about 0.5 percent by weight; and
- 30 (f) the branched carboxylic acid is selected from the group consisting of isostearic acid, 2-ethyl-hexanoic acid and mixtures thereof.

-35 -

21. The soap bar of claim 20 wherein the carboxylic acids from which the carboxylic acid salts in the soap bar are derived comprises from zero to less than about 0.5 percent by weight of C6 to C10 linear, straight chain carboxylic acids.

22. The soap bar of claim 20 wherein:

- (a) the reducing agent is sodium hydrosulfite, present at a concentration of about 0.4 percent by weight;
- 10 (b) about 50 to about 55 percent by weight of the soap bar comprises a mixture of sodium and triethanolamine salts of C6 to C22 carboxylic acids, the weight ratio of said sodium salts to said triethanolamine salts being from about 45:55 to about 55:45;
- 15 (c) at least about 90 percent by weight of the carboxylic acids from which the carboxylic acid salts in the soap bar are derived are C12 to C18 linear, straight chain saturated carboxylic acids;
- (d) about 40 to about 45 percent by weight of the soap bar comprises the solvent mixture, said solvent mixture comprising from about 70 to about 75 percent by weight triethanolamine;
- 20 (e) butylated hydroxy toluene is present in the soap bar at a concentration of about 0.01 percent by weight and
- 25 Vitamin E is present in the soap bar at a concentration of about 0.1 percent by weight; and
- (f) the branched carboxylic acid is selected from the group consisting of isostearic acid, 2-ethyl-hexanoic acid and mixtures thereof, and comprises from about 2 to
- 30 about 10 percent by weight of the total of the carboxylic acids from which the soap bar is derived.

23. The soap bar of claim 22 wherein the carboxylic acids from which the carboxylic acid salts in the soap

-36 -

bar are derived comprises from zero to less than about 0.5 percent by weight of C6 to C10 linear, straight chain carboxylic acids.

- 5 24. The soap bar consisting essentially of :

Component	Composition (percent by weight)
sodium soap	27.4
triethanolamine soap	24.7
water	12.0
triethanolamine	32
sodium hydrosulfite	0.4
BHT	0.01
Glycerine	1.0
mineral oil	0.5
coco diethanolamide	1.0
pentasodium-diethylenetriaminepentaacetate	0.5
Vitamin E	0.1

- wherein the sodium soap and triethanolamine soap are sodium and triethanolamine salts of fatty carboxylic acids; at least 90 percent of the carboxylic acids from which the carboxylic acid salts are derived are C12 to C18 linear, straight chain, saturated carboxylic acids; from about 2 to about 10 percent by weight of the carboxylic acid salts are derived from branched acids selected from 2-ethyl hexanoic acid and isostearic acid; and wherein the iodine number of the carboxylic acids from which the carboxylic acids salts in the soap bar are derived have an iodine number less than or equal to about 2.0.

-37 -

25. The soap bar of claim 24 wherein the carboxylic acids from which the carboxylic acid salts in the soap bar are derived comprises from zero to less than about 0.5 percent by weight of C6 to C10 linear, straight chain carboxylic acids.

26. A method of making a clear, colorless, soap bar which is resistant to color degradation and which is extremely mild to the skin comprising:

- 10 (a) providing a mixture of C6 to C22 carboxylic acids wherein about 2 to about 20 weight percent of the carboxylic acids are C8 to C18 branched carboxylic acids and wherein the carboxylic acids have an iodine number less than or equal to about 2.0.;
- 15 (b) heating the mixture of C6 to C22 carboxylic acids to produce a molten mixture of carboxylic acids;
- (c) admixing a reducing agent with the molten carboxylic acids;
- (d) admixing less than a molar equivalent of alkali
- 20 metal hydroxide with the C6 to C22 carboxylic acids, affording a mixture of unneutralized carboxylic acids with neutralized carboxylic acids;
- (e) admixing alkanolamine with the mixture of unneutralized carboxylic acids and neutralized
- 25 carboxylic acids, affording a mixture of alkali metal and alkanolamine salts of the C6 to C22 carboxylic acids;
- (f) providing sufficient excess of alkanolamine to function as a solvent which increases the clarity of the
- 30 soap bar, thereby providing a mixture of carboxylic acid salts and solvent comprising alkanolamine;
- (g) admixing a discoloration-retarding effective amount of an antioxidant to the carboxylic acid salts and solvent;

-38 -

(h) transferring the molten mixture of salts and solvent into molds; and

(i) cooling the molds to harden the molten soap into solid soap.

5

27. The process of claim 26 wherein steps d and e are carried out simultaneously.

28. The process of claim 26 wherein step e is carried  
10 out before step d.



1/2




Fig. 1

SUBSTITUTE SHEET (RULE 26)

2/2




Fig. 2

## INTERNATIONAL SEARCH REPORT

Internat Application No

PCT/US 97/11036

A. CLASSIFICATION OF SUBJECT MATTER  
 IPC 6 C11D9/04 C11D9/00 C11D17/00 C11D9/22

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C11D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 3 793 214 A (O NEILL J ET AL) 19 February 1974 cited in the application see the whole document ---	1-6, 11-13
A	US 2 921 907 A (JOHN G. KLEYN) 19 January 1960 see column 1, line 41 - line 46 ---	1-3
A	US 4 468 338 A (LINDBERG GORDON A) 28 August 1984 cited in the application see the whole document ---	1-5, 11-19
A	US 4 758 370 A (JUNGERMANN ERIC ET AL) 19 July 1988 cited in the application see examples ---	1-5, 11-19
-/-		

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

## \* Special categories of cited documents:

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

\*S\* document member of the same patent family

Date of the actual completion of the international search

27 October 1997

Date of mailing of the international search report

11.11.97

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
 NL - 2280 HV Rijswijk  
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
 Fax: (+31-70) 340-3016

Authorized officer

Pelli Wablat, B

## INTERNATIONAL SEARCH REPORT

Intern. Appl. Application No

PCT/US 97/11036

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP 0 335 026 A (UNILEVER PLC ET AL.) 4 October 1989 cited in the application see claims ---	1,7
A	DATABASE WPI Section Ch, Week 9433 Derwent Publications Ltd., London, GB; Class D25, AN 94-269714 XP002044691 & JP 06 200 298 A (TAMANOHADA SEKKEI KK) , 19 July 1994 see abstract ---	1
A	DATABASE WPI Section Ch, Week 9250 Derwent Publications Ltd., London, GB; Class D21, AN 92-413932 XP002044692 & SU 1 708 827 A (CHELOVEK COOP) , 30 January 1992 see abstract -----	1

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 97/11036

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 3793214 A	19-02-74	US 3926828 A	16-12-75
US 2921907 A	19-01-60	NONE	
US 4468338 A	28-08-84	NONE	
US 4758370 A	19-07-88	AU 603853 B	29-11-90
		AU 8310487 A	03-11-88
		CA 1309637 A	03-11-92
		DE 3870713 A	11-06-92
		DK 229988 A	31-10-88
		EP 0294010 A	07-12-88
		JP 1912201 C	09-03-95
		JP 6037638 B	18-05-94
		JP 63275700 A	14-11-88
EP 0335026 A	04-10-89	AU 600566 B	16-08-90
		AU 1412188 A	05-10-89

**THIS PAGE BLANK (USPTO)**